

A NOTE ON FRACTIONAL ASKEY–WILSON INTEGRALS

JIAN CAO AND SAMA ARJIKA

ABSTRACT. In this paper, we generalize fractional q -integrals by the method of q -difference equation. In addition, we deduce fractional Askey–Wilson integral, reversal type fractional Askey–Wilson integral and Ramanujan type fractional Askey–Wilson integral.

1. INTRODUCTION

The fractional q -calculus theories have been applied successfully in many fields, which is a very suitable tool in describing and solving a lot of problems in numerous sciences [23, 18], such as high-energy physics, system control, biomedical engineering and economics. Its theoretical and applied research has become a hot spot in the world. The treatment from the point view of the q -calculus can open new perspectives as it did, for example, in optimal control problems [7, 8, 15, 27]. For more information, see details in [10, 11, 2, 3, 21, 29, 30, 31, 32, 33, 24, 25, 26].

In this paper, we follow the notations and terminology in [16] and suppose that $0 < q < 1$. We first show a list of various definitions and notations in q -calculus which are useful to understand the subject of this paper. The basic hypergeometric series ${}_r\phi_s$ [16]

$${}_r\phi_s \left[\begin{matrix} a_1, a_2, \dots, a_r \\ b_1, b_2, \dots, b_s \end{matrix} ; q, z \right] = \sum_{n=0}^{\infty} \frac{(a_1, a_2, \dots, a_r; q)_n}{(q, b_1, b_2, \dots, b_s; q)_n} \left[(-1)^n q^{\binom{n}{2}} \right]^{1+s-r} z^n \quad (1)$$

converges absolutely for all z if $r \leq s$ and for $|z| < 1$ if $r = s + 1$ and for terminating. The compact factorials of ${}_r\phi_s$ are defined respectively by

$$(a; q)_0 = 1, \quad [a]_q := \frac{1 - q^a}{1 - q}, \quad (a; q)_n = \prod_{k=0}^{n-1} (1 - aq^k), \quad (a; q)_\infty = \prod_{k=0}^{\infty} (1 - aq^k) \quad (2)$$

and $(a_1, a_2, \dots, a_m; q)_n = (a_1; q)_n (a_2; q)_n \cdots (a_m; q)_n$, where $m \in \mathbb{N} := \{1, 2, 3, \dots\}$ and $n \in \mathbb{N}_0 := \mathbb{N} \cup \{0\}$. The q -gamma function is defined by [16]

$$\Gamma_q(x) = \frac{(q; q)_\infty}{(q^x; q)_\infty} (1 - q)^{1-x}, \quad x \in \mathbb{R} \setminus \{0, -1, -2, \dots\}. \quad (3)$$

The Thomae–Jackson q -integral is defined by [16, 17, 28]

2010 *Mathematics Subject Classification.* 05A30, 11B65, 33D15, 33D45, 33D60, 39A13, 39B32.
Key words and phrases. fractional q -integral; Askey–Wilson integral; q -difference equation.
Submitted Sep. 18, 2020.

$$\int_a^b f(x) d_q x = (1-q) \sum_{n=0}^{\infty} [bf(bq^n) - af(aq^n)] q^n. \quad (4)$$

The Riemann–Liouville fractional q -integral operator is introduced in [1]

$$(I_q^\alpha f)(x) = \frac{x^{\alpha-1}}{\Gamma_q(\alpha)} \int_0^x (qt/x; q)_{\alpha-1} f(t) d_q t. \quad (5)$$

The generalized Riemann–Liouville fractional q -integral operator is given by [25]

$$(I_{q,a}^\alpha f)(x) = \frac{x^{\alpha-1}}{\Gamma_q(\alpha)} \int_a^x (qt/x; q)_{\alpha-1} f(t) d_q t, \quad \alpha \in \mathbb{R}^+. \quad (6)$$

In fact, we rewrite fractional q -integral (6) equivalently as follows by (4)

$$(I_{q,a}^\alpha f)(x) = \frac{x^{\alpha-1}(1-q)}{\Gamma_q(\alpha)} \sum_{n=0}^{\infty} [x(q^{n+1}; q)_{\alpha-1} f(xq^n) - a(aq^{n+1}/x; q)_{\alpha-1} f(aq^n)] q^n.$$

In paper [13], author built the relations between the following fractional q -integrals and certain generating functions for q -polynomials.

Proposition 1 ([13, Theorem 3]). *For $\alpha \in \mathbb{R}^+$ and $0 < a < x < 1$, if $\max\{|at|, |az|\} < 1$, we have*

$$\begin{aligned} I_{q,a}^\alpha \left\{ \frac{(bxz, xt; q)_\infty}{(xs, xz; q)_\infty} \right\} \\ = \frac{(1-q)^\alpha (abz, at; q)_\infty}{(as, az; q)_\infty} \sum_{k=0}^{\infty} \frac{x^{\alpha+k} (a/x; q)_{\alpha+k}}{a^k (q; q)_{\alpha+k}} {}_3\phi_2 \left[\begin{matrix} q^{-k}, as, az \\ at, abz \end{matrix}; q, q \right]. \end{aligned} \quad (7)$$

In this paper, we generalize fractional q -integrals and give applications of fractional Askey–Wilson integrals as follows.

Theorem 2. *For $\alpha \in \mathbb{R}^+$ and $0 < a < x < 1$, if $\max\{|at|, |az|, |aru|\} < 1$, we have*

$$\begin{aligned} I_{q,a}^\alpha \left\{ \frac{(bxz, xt, xru; q)_\infty}{(xs, xz, xu; q)_\infty} \right\} \\ = \frac{(1-q)^\alpha (abz, at, aru; q)_\infty}{(as, az, au; q)_\infty} \sum_{k=0}^{\infty} \frac{x^{\alpha+k} (a/x; q)_{\alpha+k}}{a^k (q; q)_{\alpha+k}} {}_4\phi_3 \left[\begin{matrix} q^{-k}, as, az, au \\ abz, at, aru \end{matrix}; q, q \right]. \end{aligned} \quad (8)$$

Remark 3. *For $u = 0$ in Theorem 2, equation (8) reduces to (7).*

Before the proof of Theorem 2, the following lemmas are necessary.

Lemma 4 ([22, Proposition 1.2]). *Let $f(a, b, c)$ be a three-variable analytic function in a neighborhood of $(a, b, c) = (0, 0, 0) \in C^3$. If $f(a, b, c)$ satisfies the difference equation*

$$(c-b)f(a, b, c) = abf(a, bq, cq) - bf(a, b, cq) + (c-ab)f(a, bq, c), \quad (9)$$

then we have

$$f(a, b, c) = T(a, bD_c)\{f(a, 0, c)\}, \quad (10)$$

where

$$T(a, bD_c) = \sum_{n=0}^{\infty} \frac{(a; q)_n}{(q; q)_n} (bD_c)^n, \quad D_c\{f(c)\} = \frac{f(c) - f(cq)}{c}. \quad (11)$$

Lemma 5. For $\max\{|as|, |az|, |au|\} < 1$, we have

$$\begin{aligned} & (s - u) \frac{(abz, at, aru; q)_\infty}{(as, az, au)_\infty} \\ &= ur \frac{(abz, at, aruq; q)_\infty}{(asq, az, auq)_\infty} - u \frac{(abz, at, aru; q)_\infty}{(asq, az, au)_\infty} + (s - ur) \frac{(abz, at, aruq; q)_\infty}{(as, az, auq)_\infty}. \end{aligned} \tag{12}$$

Proof of Lemma 5. The right-hand side (RHS) of equation (12) equals

$$\begin{aligned} & ur \frac{(1 - xs)(1 - xu)}{1 - xur} \frac{(abz, at, aru; q)_\infty}{(as, az, au)_\infty} - u(1 - xs) \frac{(abz, at, aru; q)_\infty}{(as, az, au)_\infty} \\ & \quad + (s - ur) \frac{1 - xu}{1 - xur} \frac{(abz, at, aru; q)_\infty}{(as, az, au)_\infty} = (s - u) \frac{(abz, at, aru; q)_\infty}{(as, az, au)_\infty}, \end{aligned} \tag{13}$$

which is equals to the left-hand side (LHS) of the equation (12). The proof is complete.

Lemma 6 ([14, Eq. (2.3)]). For $\max\{|bt|, |ct|\} < 1$, we have

$$T(a, bD_c) \left\{ \frac{1}{(ct; q)_\infty} \right\} = \frac{(abt; q)_\infty}{(bt, ct; q)_\infty}. \tag{14}$$

Proof of Theorem 2. Denoting the RHS of the equation (8) by $f(r, u, s)$, and rewriting $f(r, u, s)$ equivalently by

$$f(r, u, s) = \sum_{k=0}^{\infty} \frac{x^{\alpha+k} (a/x; q)_{\alpha+k}}{a^k (q; q)_{\alpha+k}} \sum_{j=0}^k \frac{(q^{-k}; q)_j q^j}{(q; q)_j} \cdot \frac{(1 - q)^\alpha (abzq^j, atq^j, aruq^j; q)_\infty}{(asq^j, azq^j, auq^j; q)_\infty},$$

we check that $f(r, u, s)$ satisfies the equation (9) by Lemma 5, then we have

$$\begin{aligned} f(r, u, s) &= T(r, uD_s) \{f(r, 0, s)\} \\ &= T(r, uD_s) \left\{ \sum_{k=0}^{\infty} \frac{x^{\alpha+k} (a/x; q)_{\alpha+k}}{a^k (q; q)_{\alpha+k}} \sum_{j=0}^k \frac{(q^{-k}; q)_j q^j}{(q; q)_j} \cdot \frac{(1 - q)^\alpha (abzq^j, atq^j; q)_\infty}{(asq^j, azq^j; q)_\infty} \right\} \\ &= T(r, uD_s) \left\{ I_{q,a}^\alpha \left\{ \frac{(xbz, xt; q)_\infty}{(xs, xz; q)_\infty} \right\} \right\} \\ &= I_{q,a}^\alpha \left\{ T(r, uD_s) \left\{ \frac{(xbz, xt; q)_\infty}{(xs, xz; q)_\infty} \right\} \right\} \\ &= I_{q,a}^\alpha \left\{ \frac{(xbz, xt; q)_\infty}{(xz; q)_\infty} \cdot T(r, uD_s) \left\{ \frac{1}{(xs; q)_\infty} \right\} \right\}, \end{aligned}$$

which equals the LHS of the equation (8) by Lemma 6. The proof is complete.

The rest of the paper is organized as follows. In Section 2, we give the fractional Askey–Wilson integral. In Section 3, we obtain the reversal type fractional Askey–Wilson integral. In Section 4, we get the Ramanujan type fractional Askey–Wilson integral.

2. A GENERALIZATION OF ASKEY–WILSON INTEGRALS

In 1985, Askey and Wilson gave the famous Askey–Wilson integral, which greatly promoted the research and development of orthogonal polynomials. Chen and Gu [14]??Liu [19] and Cao [9] et al have promoted Askey–Wilson integral by different methods. For more information, see details in [4, 9, 14, 19].

In this section, we use the fractional q -integrals to generalize Askey–Wilson integrals.

Proposition 7 ([4, Theorem 2.1]). *If $\max\{|a|, |b|, |c|, |d|\} < 1$, we have*

$$\int_0^\pi \frac{h(\cos 2\theta; 1)}{h(\cos \theta; a, b, c, d)} d\theta = \frac{2\pi(abcd; q)_\infty}{(q, ab, ac, ad, bc, bd, cd; q)_\infty}, \tag{15}$$

where

$$h(\cos \theta; a) = (ae^{i\theta}, ae^{-i\theta}; q)_\infty,$$

$$h(\cos \theta; a_1, a_2, \dots, a_m) = h(\cos \theta; a_1)h(\cos \theta; a_2) \cdots h(\cos \theta; a_m).$$

Theorem 8. *For $\alpha \in \mathbb{R}^+$, if $\max\{|a|, |b|, |c|, |d|\} < 1$, we have*

$$\int_0^\pi \frac{h(\cos 2\theta; 1)}{h(\cos \theta; a, b, c, d)} \sum_{k=0}^\infty \frac{x^{\alpha+k}(a/x; q)_{\alpha+k}}{a^k(q; q)_{\alpha+k}} {}_4\phi_3 \left[\begin{matrix} q^{-k}, abcd, ae^{i\theta}, ae^{-i\theta} \\ ab, ac, ad \end{matrix}; q, q \right] d\theta$$

$$= \frac{2\pi(abcd; q)_\infty}{(q, ab, ac, ad, bc, bd, cd; q)_\infty} \frac{x^\alpha(a/x; q)_\alpha}{(q; q)_\alpha}. \tag{16}$$

Corollary 9. *For $\alpha \in \mathbb{R}^+$, if $\max\{|a|, |b|, |c|\} < 1$, we have*

$$\int_0^\pi \frac{h(\cos 2\theta; 1)}{h(\cos \theta; a, b, c)} \sum_{k=0}^\infty \frac{x^{\alpha+k}(a/x; q)_{\alpha+k}}{a^k(q; q)_{\alpha+k}} {}_3\phi_2 \left[\begin{matrix} q^{-k}, ae^{i\theta}, ae^{-i\theta} \\ ab, ac \end{matrix}; q, q \right] d\theta$$

$$= \frac{2\pi}{(q, ab, ac, bc; q)_\infty} \frac{x^\alpha(a/x; q)_\alpha}{(q; q)_\alpha}. \tag{17}$$

Remark 10. *For $d = 0$ in Theorem 8, equation (16) reduces to (17).*

Proof of Theorem 8. The equation (16) can be rewrite equivalently by

$$\int_0^\pi \frac{h(\cos 2\theta; 1)}{h(\cos \theta; b, c, d)} \frac{(ab, ac, ad; q)_\infty}{(ae^{i\theta}, ae^{-i\theta}, abcd; q)_\infty} \sum_{k=0}^\infty \frac{x^{\alpha+k}(a/x; q)_{\alpha+k}}{a^k(q; q)_{\alpha+k}}$$

$$\times {}_4\phi_3 \left[\begin{matrix} q^{-k}, abcd, ae^{i\theta}, ae^{-i\theta} \\ ab, ac, ad \end{matrix}; q, q \right] d\theta = \frac{2\pi}{(q, bc, bd, cd; q)_\infty} \frac{x^\alpha(a/x; q)_\alpha}{(q; q)_\alpha}. \tag{18}$$

By using Theorem 2, the LHS of the equation (18) is equivalent to

$$\int_0^\pi \frac{h(\cos 2\theta; 1)}{h(\cos \theta; b, c, d)} I_{q,a}^\alpha \left\{ \frac{(xb, xc, xd; q)_\infty}{(xe^{i\theta}, xe^{-i\theta}, xbcd; q)_\infty} \right\} d\theta$$

$$= I_{q,a}^\alpha \left\{ \int_0^\pi \frac{h(\cos 2\theta; 1)(xb, xc, xd; q)_\infty}{h(\cos \theta; x, b, c, d)(xbcd; q)_\infty} d\theta \right\}$$

$$= I_{q,a}^\alpha \left\{ \frac{2\pi}{(q, bc, bd, cd; q)_\infty} \right\}$$

$$= \frac{2\pi}{(q, bc, bd, cd; q)_\infty} \frac{x^\alpha(a/x; q)_\alpha}{(q; q)_\alpha},$$

which is the RHS of the equation (18). The proof is complete.

3. A GENERALIZATION OF REVERSAL TYPE ASKEY–WILSON INTEGRALS

In this section, we use the fractional q -integrals to expand reversal Askey–Wilson integrals.

Proposition 11 (Reversal Askey–Wilson integral [5]). *For $|qabcd| < 1$, there holds*

$$\int_{-\infty}^{\infty} \frac{h(i \sinh x; qa, qb, qc, qd)}{h(\cosh 2x; -q)} dx = \frac{(q, qab, qac, qad, qbc, qbd, qcd; q)_{\infty}}{(qabcd; q)_{\infty}} \log(q^{-1}), \tag{19}$$

where

$$h(i \sinh \alpha x; t) = \prod_{k=0}^{\infty} (1 - 2iq^k t \sinh \alpha x + q^{2k} t^2) = (ite^{\alpha x}, -ite^{-\alpha x}; q)_{\infty}.$$

Theorem 12. *For $\alpha \in \mathbb{R}^+$ and $|qabcd| < 1$, we have*

$$\begin{aligned} & \int_{-\infty}^{\infty} \frac{h(i \sinh t; qa, qb, qc, qd)}{h(\cosh 2t; -q)} \sum_{k=0}^{\infty} \frac{x^{\alpha+k} (a/x; q)_{\alpha+k}}{a^k (q; q)_{\alpha+k}} {}_4\phi_3 \left[\begin{matrix} q^{-k}, qab, qac, qad \\ iaqe^t, -iaqe^{-t}, qabcd \end{matrix}; q, q \right] dt \\ &= \frac{(q, qab, qac, qad, qbc, qbd, qcd; q)_{\infty}}{(qabcd; q)_{\infty}} \frac{x^{\alpha} (a/x; q)_{\alpha}}{(q; q)_{\alpha}} \log(q^{-1}). \end{aligned} \tag{20}$$

Corollary 13. *For $\alpha \in \mathbb{R}^+$, we have*

$$\begin{aligned} & \int_{-\infty}^{\infty} \frac{h(i \sinh t; qa, qb, qc)}{h(\cosh 2t; -q)} \sum_{k=0}^{\infty} \frac{x^{\alpha+k} (a/x; q)_{\alpha+k}}{a^k (q; q)_{\alpha+k}} {}_3\phi_2 \left[\begin{matrix} q^{-k}, qab, qac \\ iaqe^t, -iaqe^{-t} \end{matrix}; q, q \right] dt \\ &= \frac{(q, qab, qac, qbc; q)_{\infty} x^{\alpha} (a/x; q)_{\alpha}}{(q; q)_{\alpha}} \log(q^{-1}). \end{aligned} \tag{21}$$

Remark 14. *For $d = 0$ in Theorem 12, equation (20) reduces to (19).*

Proof of Theorem 12. The equation (20) can be rewrite

$$\begin{aligned} & \int_{-\infty}^{\infty} \frac{h(i \sinh t; qb, qc, qd)}{h(\cosh 2t; -q)} \frac{(iaqe^t, -iaqe^{-t}, qabcd; q)_{\infty}}{(qab, qac, qad; q)_{\infty}} \sum_{k=0}^{\infty} \frac{x^{\alpha+k} (a/x; q)_{\alpha+k}}{a^k (q; q)_{\alpha+k}} \\ & \quad \times {}_4\phi_3 \left[\begin{matrix} q^{-k}, qab, qac, qad \\ iaqe^t, -iaqe^{-t}, qabcd \end{matrix}; q, q \right] dt \\ &= (q, qbc, qbd, qcd; q)_{\infty} \frac{x^{\alpha} (a/x; q)_{\alpha}}{(q; q)_{\alpha}} \log(q^{-1}), \end{aligned} \tag{22}$$

By using Theorem 2, the LHS of the equation (22) equals

$$\begin{aligned} & \int_{-\infty}^{\infty} \frac{h(i \sinh t; qb, qc, qd)}{h(\cosh 2t; -q)} I_{q,a}^{\alpha} \left\{ \frac{(ixqe^t, -ixqe^{-t}, qxbcd; q)_{\infty}}{(qxb, qxc, qxd; q)_{\infty}} \right\} dt \\ &= I_{q,a}^{\alpha} \left\{ \int_{-\infty}^{\infty} \frac{h(i \sinh t; qx, qb, qc, qd)(qxbcd; q)_{\infty}}{h(\cosh 2t; -q)(qxb, qxc, qxd; q)_{\infty}} dt \right\} \\ &= I_{q,a}^{\alpha} \left\{ (q, qbc, qbd, qcd; q)_{\infty} \log(q^{-1}) \right\} \\ &= (q, qbc, qbd, qcd; q)_{\infty} \frac{x^{\alpha} (a/x; q)_{\alpha}}{(q; q)_{\alpha}} \log(q^{-1}). \end{aligned}$$

which is the RHS of the equation (22). The proof is complete.

4. A GENERALIZATION OF RAMANUJAN TYPE ASKEY–WILSON INTEGRALS

In 1994, Atakishiyev discovered the Atakishiyev integral by Ramanujan’s method. After that, Wang [33] and Liu [20] et al conducted different generalization and proving lines for this integral.

In this part, we get the generating functions of Atakishiyev integral by fractional q -integral.

Proposition 15 (Atakishiyev integral [6]). *If α is a real number and $q = \exp(-2\alpha^2)$, then we have*

$$\int_{-\infty}^{\infty} h(i \sinh \alpha t; a, b, c, d) e^{-t^2} \cosh \alpha t dt = \sqrt{\pi} q^{-\frac{1}{8}} \frac{(ab/q, ac/q, ad/q, bc/q, bd/q, cd/q; q)_{\infty}}{(abcd/q^3; q)_{\infty}}. \tag{23}$$

Theorem 16. *For $\alpha \in \mathbb{R}^+$ and $|abcd/q^3| < 1$, if α is a real number and $q = \exp(-2\alpha^2)$, then we have*

$$\begin{aligned} & \int_{-\infty}^{\infty} h(i \sinh \alpha t; a, b, c, d) e^{-x^2} \cosh \alpha t \sum_{k=0}^{\infty} \frac{x^{\alpha+k} (a/x; q)_{\alpha+k}}{a^k (q; q)_{\alpha+k}} \\ & \times {}_4\phi_3 \left[\begin{matrix} q^{-k}, ab/q, ac/q, ad/q \\ iae^{\alpha t}, -iae^{-\alpha t}, abcd/q^3 \end{matrix}; q, q \right] dt \\ & = \sqrt{\pi} q^{-\frac{1}{8}} \frac{(ab/q, ac/q, ad/q, bc/q, bd/q, cd/q; q)_{\infty}}{(abcd/q^3; q)_{\infty}} \frac{x^{\alpha} (a/x; q)_{\alpha}}{(q; q)_{\alpha}}. \end{aligned} \tag{24}$$

Corollary 17. *For $\alpha \in \mathbb{R}^+$, if α is a real number and $q = \exp(-2\alpha^2)$, then we have*

$$\begin{aligned} & \int_{-\infty}^{\infty} h(i \sinh \alpha t; a, b, c) e^{-x^2} \cosh \alpha t \sum_{k=0}^{\infty} \frac{x^{\alpha+k} (a/x; q)_{\alpha+k}}{a^k (q; q)_{\alpha+k}} {}_3\phi_2 \left[\begin{matrix} q^{-k}, ab/q, ac/q \\ iae^{\alpha t}, -iae^{-\alpha t} \end{matrix}; q, q \right] dt \\ & = \sqrt{\pi} q^{-\frac{1}{8}} (ab/q, ac/q, bc/q; q)_{\infty} \frac{x^{\alpha} (a/x; q)_{\alpha}}{(q; q)_{\alpha}}. \end{aligned} \tag{25}$$

Remark 18. *For $d = 0$ in Theorem 16, equation (24) reduces to (25).*

Proof of Theorem 16. The equation (24) can be rewritten

$$\begin{aligned} & \int_{-\infty}^{\infty} h(i \sinh \alpha t; b, c, d) e^{-x^2} \cosh \alpha t \frac{(iae^{\alpha t}, -iae^{-\alpha t}, abcd/q^3; q)_{\infty}}{(ab/q, ac/q, ad/q; q)_{\infty}} \sum_{k=0}^{\infty} \frac{x^{\alpha+k} (a/x; q)_{\alpha+k}}{a^k (q; q)_{\alpha+k}} \\ & \times {}_4\phi_3 \left[\begin{matrix} q^{-k}, ab/q, ac/q, ad/q \\ iae^{\alpha t}, -iae^{-\alpha t}, abcd/q^3 \end{matrix}; q, q \right] dt \\ & = \sqrt{\pi} q^{-\frac{1}{8}} (bc/q, bd/q, cd/q; q)_{\infty} \frac{x^{\alpha} (a/x; q)_{\alpha}}{(q; q)_{\alpha}}. \end{aligned} \tag{26}$$

By using Theorem 2, the LHS of the equation (26) is equal to

$$\begin{aligned} & \int_{-\infty}^{\infty} h(i \sinh \alpha t; b, c, d) e^{-x^2} \cosh \alpha t \cdot I_{q,a}^{\alpha} \left\{ \frac{(ixe^{\alpha t}, -ixe^{-\alpha t}, xbcd/q^3; q)_{\infty}}{(xb/q, xc/q, xd/q; q)_{\infty}} \right\} dt \\ &= I_{q,a}^{\alpha} \left\{ \int_{-\infty}^{\infty} h(i \sinh \alpha t; x, b, c, d) e^{-x^2} \cosh \alpha t \cdot \frac{(xbcd/q^3; q)_{\infty}}{(xb/q, xc/q, xd/q; q)_{\infty}} dt \right\} \\ &= I_{q,a}^{\alpha} \left\{ \sqrt{\pi} q^{-\frac{1}{8}} (bc/q, bd/q, cd/q; q)_{\infty} \right\} \\ &= \sqrt{\pi} q^{-\frac{1}{8}} (bc/q, bd/q, cd/q; q)_{\infty} \frac{x^{\alpha}(a/x; q)_{\alpha}}{(q; q)_{\alpha}}, \end{aligned}$$

which is the RHS of the equation (26). The proof is complete.

ACKNOWLEDGMENTS

The author would like to thank the referees and editors for their many valuable comments and suggestions. This work was supported by the Zhejiang Provincial Natural Science Foundation of China (No. LY21A010019).

REFERENCES

- [1] W.A. Al-Salam, *Some fractional q -integral and q -derivatives*, Proc. Edinburgh Math. Soc. **15**(1966), 135–140.
- [2] W.A. Al-Salam, *q -Analogues of Cauchy's formulas*, Proc. Amer. Math. Soc. **17**(1966), 616–621.
- [3] G.E. Andrews, *Applications of basic hypergeometric series*, SIAM Rev. **16**(1974), 441–484.
- [4] R. Askey, J.A. Willson, *Some basic hypergeometric orthogonal polynomials that generalize Jacobi polynomials*, Mem. Amer. Math. Soc. **319**(1985) 55.
- [5] R. Askey, *Bate integrals and q -extensions*, in: Proceedings of the Ramanujan Centennial International Conference, Anna-malainagar, 15-18 December 1987, pp. 85-102.
- [6] N.M. Atakishiyev, *On the Askey–Wilson q -beta integral*, Theoretical and Mathematical Physics, **99**(1994), 155–159.
- [7] D. Baleanu, A. Fernandez, *On some new properties of fractional derivatives with Mittag-Leffler kernel*, Commun Nonlinear Sci Numer Simulat. **59**(2018), 444–462.
- [8] G. Bangerezako, *Variational calculus on q -nonuniform lattices*, J. Math. Anal. Appl. **306**(2005), 161–179.
- [9] J. Cao, *A note on generalized q -difference equations for q -beta and Andrews–Askey integral*, J. Math. Anal. Appl. **412**(2014), 841–851.
- [10] J. Cao and D.-W. Niu, *q -Difference equations for Askey–Wilson type integrals via q -polynomials*, J. Math. Anal. Appl. **452**(2017), 830–845.
- [11] J. Cao, H.M. Srivastava, Z.G. Liu, *Some iterated fractional q -integrals and their applications*, Fractional Calculus and Applied Analysis. **21**(2018), 672–695.
- [12] J. Cao, *A note on q -difference equations for Ramanujan's integrals*, The Ramanujan Journal. (2018), 1–11.
- [13] J. Cao, *A note on fractional q -integrals and applications to generating functions and q -Mittag–Leffler functions*, J. Frac. Calc. Appl. **10**(2019), 136–146.
- [14] V.Y.B. Chen, S.S. Gu, *The Cauchy operator for basic hypergeometric series*, Adv. Appl. Math. **41**(2008), 177–196.
- [15] M. El-Shahed, M. Gaber and M. Al-Yami, *The fractional q -differential transformation and its application*, Commun Nonlinear Sci Numer Simulat. **18**(2013), 42–55.
- [16] G. Gasper, M. Rahman, *Basic Hypergeometric Series*, Camberdge: Camberdge Univ Press, 2004.
- [17] F.H. Jackson, *On q -definite integrals*, Q. J. Pure Appl. Math. **41**(1910), 193–203.
- [18] A.A. Kilbas, H.M. Srivastava and J.J. Trujillo, *Theory and Applications of Fractional Differential Equations*. North-Holland Math. Studies, Vol. 204, Elsevier(North-Holland)Sci.Publ, Amsterdam,London and New York,2006.

- [19] Z.-G. Liu, *An identity of Andrews and the Askey–Wilson integral*, J. Ramanujan J. **19**(2009), 115–119.
- [20] Z.-G. Liu, *Two expansion formulas involving the Rogers–Szegő polynomials with applications*, J. International Journal of Number Theory **11**(2015), 507–525.
- [21] Z.-G. Liu, *On a reduction formula for a kind of double q -integrals identities*, Symmetry **8**(2016), Article ID 44, 1–16.
- [22] D.-Q. Lu, *q -difference equation and cauchy operator identities*, J. Math. Anal. Appl. **359**(2009), 265–274.
- [23] I. Podlubny, *Fractional Differential Equations, An Introduction to Fractional Derivatives, Fractional Differential Equations, Some Methods of Their Solution and Some of Their Applications*, Academic Press, San Diego–Boston–New York–London–Tokyo–Toronto, 1999.
- [24] Predrag M. Rajković, Sladjana D. Marinković and Miomir S. Stanković, *On q -analogues of Caputo derivative and Mittag–Leffler function*, Fract. Calc. Appl. Anal. **10**(2007), 359–373.
- [25] Predrag M. Rajković, Sladjana D. Marinković and Miomir S. Stanković, *Fractional integrals and derivatives in q -calculus*, Appl. Anal. Discrete Math. **1**(2007), 311–323.
- [26] Predrag M. Rajković, Sladjana D. Marinković and Miomir S. Stanković, *A generalization of the concept of q -fractional integrals*, Acta Math. Sin., Engl. Ser., **25**(2009), 1635–1646.
- [27] K. Sayevand, J.T. Machado and D. Baleanu, *A new glance on the Leibniz rule for fractional derivatives*, Commun Nonlinear Sci Numer Simulat. **62**(2018), 244–249.
- [28] J. Thomae, *Beiträge zur Theorie der durch die Heinesche Reihe: Darstellbaren Function*, J. Reine Angew. Math. **70**(1869), 258–281.
- [29] M. Wang, *A recurring q -integral formula*, Appl. Math. Lett. **23**(2009), 256–260.
- [30] M. Wang, *A remark on Andrews–Askey integral*, J. Math. Anal. Appl. **341**(2008), 1487–1494.
- [31] M. Wang, *Generalizations of Milne’s $U(n + 1)$ q -binomial theorem*, Comput. Math. Appl. **58**(2009), 80–87.
- [32] M. Wang, *q -integral representation of the Al-Salam–Carlitz polynomials*, Appl. Math. Lett. **22**(2009), 943–945.
- [33] M. Wang, *An extension of q -beta integral with application*, J. Math. Anal. Appl. **365**(2010), 653–658.

JIAN CAO

DEPARTMENT OF MATHEMATICS, HANGZHOU NORMAL UNIVERSITY, HANGZHOU CITY, ZHEJIANG PROVINCE, 311121, PR CHINA

E-mail address: 21caojian@163.com, 21caojian@hznu.edu.cn

SAMA ARJIKA

DEPARTMENT OF MATHEMATICS AND INFORMATICS, UNIVERSITY OF AGADEZ, NIGER

E-mail address: rjksama2008@gmail.com